

ON COMPONENT GROUPS OF JACOBIANS OF QUATERNIONIC MODULAR CURVES

MIHRAN PAPIKIAN

Dedicated to Ernst-Ulrich Gekeler

ABSTRACT. We use a combinatorial result relating the discriminant of the cycle pairing on a weighted finite graph to the eigenvalues of its Laplacian, to deduce a formula for the orders of component groups of Jacobians of modular curves arising from quaternion algebras over $\mathbb{F}_q(T)$ or \mathbb{Q} . Our formula over \mathbb{Q} recovers a result of Jordan and Livné.

1. INTRODUCTION

Let \mathbb{F}_q be a finite field with q elements, where q is a power of a prime number p . Let $A = \mathbb{F}_q[T]$ be the ring of polynomials in the indeterminate T with coefficients in \mathbb{F}_q , and $F = \mathbb{F}_q(T)$ be the fraction field of A . The degree map $\deg : F \rightarrow \mathbb{Z} \cup \{-\infty\}$, which assigns to a non-zero polynomial its degree in T and $\deg(0) = -\infty$, is a valuation of F . The corresponding place of F will be denoted by ∞ . Apart from ∞ , the places of F are in bijection with the non-zero prime ideals of A . Given a place v of F , we denote by F_v the completion of F at v , and by \mathbb{F}_v the residue field of F_v .

Let \mathcal{B} be a division quaternion algebra over F , which is “indefinite” in the sense that $\mathcal{B} \otimes_F F_\infty \cong \mathbb{M}_2(F_\infty)$. Let \mathfrak{d} be the (reduced) discriminant of \mathcal{B} ; cf. [20, p. 58]. This is the product of primes $\mathfrak{p} \triangleleft A$ where \mathcal{B} ramifies, i.e., $\mathcal{B} \otimes_F F_{\mathfrak{p}}$ is a division algebra over $F_{\mathfrak{p}}$. It is known from the theory of quaternion algebras that \mathfrak{d} is a product of an even number of primes, and, up to isomorphism, \mathfrak{d} uniquely determines \mathcal{B} ; cf. [20]. (Moreover, any square-free \mathfrak{d} with an even number of prime divisors is the discriminant of some indefinite quaternion algebra over F .)

There is a smooth projective curve $X^{\mathfrak{d}}$ over F associated with the quaternion algebra \mathcal{B} . It is the coarse moduli scheme of the so-called \mathcal{D} -elliptic sheaves; roughly, a \mathcal{D} -elliptic sheaf is a vector bundle equipped with a meromorphic Frobenius and an action of \mathcal{B} ; we refer to [10] for the precise definitions. The curve $X^{\mathfrak{d}}$ has bad reduction exactly at the primes dividing \mathfrak{d} and ∞ ; cf. [10]. Let $J^{\mathfrak{d}}$ be the Jacobian variety of $X^{\mathfrak{d}}$, and $\mathcal{J}^{\mathfrak{d}}$ be the Néron model of $J^{\mathfrak{d}}$ over A . Let \mathfrak{p} be a prime dividing \mathfrak{d} . The main result of this paper is a formula for the order of the component group $\Phi_{J^{\mathfrak{d}}, \mathfrak{p}} = (\mathcal{J}^{\mathfrak{d}} \otimes_{\mathbb{F}_{\mathfrak{p}}} \overline{\mathbb{F}_{\mathfrak{p}}}) / (\mathcal{J}^{\mathfrak{d}} \otimes_{\mathbb{F}_{\mathfrak{p}}} \mathbb{F}_{\mathfrak{p}})^0$. In order to state this formula, we need to introduce more notation.

The order of a finite set S will be denoted by $|S|$. We define norm and degree for a non-zero ideal \mathfrak{n} of A by $|\mathfrak{n}| := |A/\mathfrak{n}|$ and $\deg(\mathfrak{n}) := \log_q |\mathfrak{n}|$. The prime ideals

2010 *Mathematics Subject Classification.* 11G18, 05C25, 11F12.

Key words and phrases. Graph Laplacian; Component group; Shimura curve.

The author’s research was partially supported by grants from the Simons Foundation (245676) and the National Security Agency (H98230-15-1-0008).

of A will always be assumed to be non-zero. Let \mathcal{B}' be the “definite” quaternion algebra over F which is ramified at ∞ and at the primes $\mathfrak{p}_1, \dots, \mathfrak{p}_s$ appearing in the prime decomposition of

$$\mathfrak{d}' := \mathfrak{d}/\mathfrak{p} = \prod_{i=1}^s \mathfrak{p}_i.$$

Fix a maximal A -order \mathcal{M} in \mathcal{B}' . Let $h(\mathfrak{d}')$ be the class number of \mathcal{B}' , that is, the number of left \mathcal{M} ideal classes. It is known that $h(\mathfrak{d}')$ is finite and independent of the choice of \mathcal{M} . Let $I_1, I_2, \dots, I_{h(\mathfrak{d}')}$ be representatives of the left ideal classes of \mathcal{M} , and let \mathcal{M}_i be the right order of I_i , $1 \leq i \leq h(\mathfrak{d}')$. It is not hard to show that each \mathcal{M}_i^\times is isomorphic either to \mathbb{F}_q^\times or $\mathbb{F}_{q^2}^\times$; thus, $w_i := |\mathcal{M}_i^\times/A^\times|$ is either equal to 1 or $q+1$; see [3, p. 383]. It is known that (cf. [3, (1)])

$$m(\mathfrak{d}') := \sum_{i=1}^{h(\mathfrak{d}')} w_i^{-1} = \frac{1}{q^2-1} \prod_{i=1}^s (|\mathfrak{p}_i| - 1);$$

$m(\mathfrak{d}')$ is called the *mass* of \mathcal{B}' . If we denote by h_{q+1} the number of w_i 's equal to $q+1$, then by (1), (4) and (6) in [3]:

$$h_{q+1} = \frac{1}{2} \prod_{i=1}^s \left(1 - (-1)^{\deg(\mathfrak{p}_i)}\right).$$

This gives a formula for the class number:

$$h(\mathfrak{d}') = m(\mathfrak{d}') + h_{q+1} \frac{q}{q+1}.$$

Definition 1.1. Denote $M_{ij} := I_j^{-1}I_i$. Let m_{ij} be a generator of the ideal $\{\text{Nr}(\alpha) \mid \alpha \in M_{ij}\} \triangleleft A$, where $\text{Nr} : \mathcal{B}' \rightarrow F$ is the reduced norm. Let $\mathfrak{m} \triangleleft A$ be a non-zero ideal. Let

$$b_{ij}(\mathfrak{m}) = \frac{\#\{\beta \in M_{ij} \mid (\text{Nr}(\beta)/m_{ij}) = \mathfrak{m}\}}{(q-1)w_j}.$$

The \mathfrak{m} -th Brandt matrix of \mathcal{B}' is $B(\mathfrak{m}) = (b_{ij}(\mathfrak{m}))_{1 \leq i, j \leq h(\mathfrak{d}')} \in \mathbb{M}_{h(\mathfrak{d}')}(\mathbb{Z})$.

Let $P(x)$ be the characteristic polynomial of the Brandt matrix $B(\mathfrak{p})$. Denote by $P'(x)$ the derivative of $P(x)$. Let

$$n(\mathfrak{d}) = \left(1 - (-1)^{\deg(\mathfrak{p})}\right) h_{q+1} = \frac{1}{2} \prod_{\substack{\mathfrak{q} \mid \mathfrak{d} \\ \mathfrak{q} \text{ is prime}}} \left(1 - (-1)^{\deg(\mathfrak{q})}\right).$$

Theorem 1.2. *We have:*

$$|\Phi_{J^\circ, \mathfrak{p}}| = \frac{|P(-|\mathfrak{p}| - 1) \cdot P'(|\mathfrak{p}| + 1)|}{2m(\mathfrak{d}')(q+1)^{n(\mathfrak{d})}}.$$

This formula is the function field analogue of a result of Jordan and Livné [8] for Shimura curves over \mathbb{Q} . In [8], for a prime p dividing the discriminant d of an indefinite quaternion algebra, the authors use the Cherednik-Drinfeld p -adic uniformization of the corresponding Shimura curve X^d to construct a regular model \mathcal{X}^d of X^d over \mathbb{Z}_p . Then, they apply a result of Raynaud (cf. [1, §9.6]), reducing the problem of computation of the group of connected components $\Phi_{J^d, p}$ of the Jacobian of X^d to linear algebra computations with the intersection matrix of the special fibre of \mathcal{X}^d . This approach leaves it somewhat mysterious the reason for

the appearance of different invariants of the quaternion algebra in the formula for $|\Phi_{J^d, \mathfrak{p}}|$.

To prove Theorem 1.2 we take a different approach, which emphasises the combinatorial nature of the formula for $|\Phi_{J^d, \mathfrak{p}}|$, although it still relies on the function field analogue of the Cherednik-Drinfeld uniformization. In Section 2, we recall a result from [13], which relates the discriminant of a weighted cycle pairing on the first homology group of a graph to the eigenvalues of its Laplacian. By a theorem of Grothendieck, given a semistable curve, the discriminant of the cycle pairing on its dual graph is the order of the component group of its Jacobian (see Theorem 2.2). On the other hand, using the rigid-analytic uniformization of $X^\mathfrak{d}$, one shows that the Laplacian of the dual graph of $X^\mathfrak{d}$ over $F_{\mathfrak{p}}$ is related to a Hecke operator with characteristic polynomial $P(x)$. Here we should mention that the dual graph of the semistable model of $X^\mathfrak{d}$ that one obtains through the rigid-analytic uniformization is naturally weighted and the Hecke operator mentioned above is related to the weighted Laplacian, so one has to deal with weighted graphs from the outset. The proof of Theorem 1.2 then reduces to computing the weights of this dual graph, which is done in Section 3. Our approach applies to Shimura curves over \mathbb{Q} as well. In Section 4, we reprove the result of Jordan and Livné using this method. (Here we also use the opportunity to correct a small mistake in the main formula of [8]; see Remark 4.1.)

Example 1.3. We observe that $h(\mathfrak{d}') = 1$ if and only if $\deg(\mathfrak{d}') \leq 2$. Since \mathfrak{d}' is a product of an odd number of primes, we conclude that $h(\mathfrak{d}') = 1$ is equivalent to $\mathfrak{d}' = \mathfrak{q}$ being a prime of degree ≤ 2 . Thus, $\mathfrak{d} = \mathfrak{p}\mathfrak{q}$ and $P(x) = x - (|\mathfrak{p}| + 1)$, since $|\mathfrak{p}| + 1$ is a root of $P(x)$ for any \mathfrak{p} (cf. [21, p. 739]). First, assume $\deg(\mathfrak{q}) = 1$. In this case, $m(\mathfrak{d}') = (q + 1)^{-1}$, and

$$n(\mathfrak{d}) = \begin{cases} 2 & \text{if } \deg(\mathfrak{p}) \text{ is odd,} \\ 0 & \text{if } \deg(\mathfrak{p}) \text{ is even.} \end{cases}$$

Thus,

$$|\Phi_{J^d, \mathfrak{p}}| = \begin{cases} \frac{|\mathfrak{p}|+1}{q+1}, & \text{if } \deg(\mathfrak{p}) \text{ is odd;} \\ (|\mathfrak{p}| + 1)(q + 1), & \text{if } \deg(\mathfrak{p}) \text{ is even.} \end{cases}$$

Now assume $\deg(\mathfrak{q}) = 2$. In this case, $m(\mathfrak{d}') = 1$, $n(\mathfrak{d}) = 0$, and

$$|\Phi_{J^d, \mathfrak{p}}| = |\mathfrak{p}| + 1.$$

Example 1.4. Some calculations of Brandt matrices over $\mathbb{F}_q(T)$ were carried out by Schweizer in [17]. For example, he computed that for $q = 2$, $\mathfrak{p} = T$, and $\mathfrak{d}' = T^5 + T^2 + 1$ we have

$$P(x) = (x-3)(x^2 + x - 1) \times (x^8 + x^7 - 11x^6 - 8x^5 + 38x^4 + 16x^3 - 44x^2 - 4x + 4).$$

Therefore, for $\mathfrak{d} = T(T^5 + T^2 + 1)$ and $\mathfrak{p} = T$, we have $h(\mathfrak{d}') = 11$, $m(\mathfrak{d}') = 31/3$, $n(\mathfrak{d}) = 2$ and

$$|\Phi_{J^d, T}| = \frac{|P(-3)P'(3)|}{2^{\frac{31}{3}}3^2} = 5^2 \cdot 11 \cdot 61 \cdot 113.$$

Nowadays, Brandt matrix calculations over $\mathbb{F}_q(T)$ are implemented in **Magma**, which makes the calculation of $|\Phi_{J^d, \mathfrak{p}}|$ fairly routine. Some of the values of $|\Phi_{J^d, \mathfrak{p}}|$ are listed in Tables 1 and 2.

\mathfrak{p}	$ \Phi_{J^\circ, \mathfrak{p}} $
$T - 3$	2^5
$T^2 - 3$	$2^{13} \cdot 3^6 \cdot 5 \cdot 7 \cdot 11 \cdot 13 \cdot 17 \cdot 37 \cdot 59$
$T^3 + T + 1$	$2^{16} \cdot 3^9 \cdot 5 \cdot 7 \cdot 11 \cdot 13 \cdot 23 \cdot 181$

TABLE 1. $q = 5$ and $\mathfrak{d} = T(T - 1)(T - 2)\mathfrak{p}$

\mathfrak{p}	$ \Phi_{J^\circ, \mathfrak{p}} $
T	$2^{10} \cdot 19^2 \cdot 43^3$
$T - 1$	$2^{12} \cdot 3 \cdot 7 \cdot 179 \cdot 1871$
$T - 3$	$2^{19} \cdot 5 \cdot 103^2$
$T^2 - 3$	$2^{23} \cdot 5^2 \cdot 7 \cdot 29 \cdot 6525499 \cdot 13235507$
$T^3 + 2$	$2^{22} \cdot 3^3 \cdot 5^6 \cdot 7 \cdot 13^4 \cdot 19^5 \cdot 43^7$

TABLE 2. $q = 7$ and $\mathfrak{d} = (T^3 - 2)\mathfrak{p}$

Let $X_0(\mathfrak{d})$ be the Drinfeld modular curve over F classifying rank-2 Drinfeld A -modules with $\Gamma_0(\mathfrak{d})$ -level structures; cf. [4]. Let $J_0(\mathfrak{d})$ be the Jacobian variety of $X_0(\mathfrak{d})$. The Jacquet-Langlands correspondence and Zarhin's isogeny theorem over function fields imply that there is a surjective homomorphism $J_0(\mathfrak{d}) \rightarrow J^\circ$ defined over F ; cf. [14, Thm. 7.1]. Thus, there are functorial homomorphisms $\psi_{\mathfrak{p}} : \Phi_{J_0(\mathfrak{d}), \mathfrak{p}} \rightarrow \Phi_{J^\circ, \mathfrak{p}}$ for primes \mathfrak{p} dividing \mathfrak{d} , where $\Phi_{J_0(\mathfrak{d}), \mathfrak{p}}$ denotes the group of connected components of the Néron model of $J_0(\mathfrak{d})$ at \mathfrak{p} . The group $\Phi_{J_0(\mathfrak{d}), \mathfrak{p}}$ is well-understood: its complete description as an abelian group can be found in [16, Thm. 5.3]. In particular, $\Phi_{J_0(\mathfrak{d}), \mathfrak{p}}$ has no elements of p -power order and the isomorphism class of $\Phi_{J_0(\mathfrak{d}), \mathfrak{p}}$ depends only on the degrees of the primes dividing \mathfrak{d} and the number of these primes. We observe from Tables 1 and 2 that these properties do not transfer to $\Phi_{J^\circ, \mathfrak{p}}$, in the sense that $|\Phi_{J^\circ, \mathfrak{p}}|$ can be divisible by the characteristic of the field and $|\Phi_{J^\circ, \mathfrak{p}}|$ depends not just on the degree of \mathfrak{p} but on the actual ideal \mathfrak{p} . Another difference between the groups $\Phi_{J_0(\mathfrak{d}), \mathfrak{p}}$ and $\Phi_{J^\circ, \mathfrak{p}}$ is that the latter is generally a much larger group. Indeed, the order of the group $\Phi_{J_0(\mathfrak{d}), \mathfrak{p}}$ grows linearly with $|\mathfrak{d}|$; see [16, Thm. 5.3]. On the other hand, one can use Theorem 1.2 and the arguments in [13] to prove the following estimates:

Theorem 1.5. *For two positive real valued functions $f(x)$ and $g(x)$ defined on an infinite subset of ideals of A , we write $f(x) \sim g(x)$ when $\lim_{|x| \rightarrow \infty} f(x)/g(x) = 1$. We have:*

- (1) *If \mathfrak{d}' is fixed and \mathfrak{p} varies, then*

$$\log_q |\Phi_{J^\circ, \mathfrak{p}}| \sim (2h(\mathfrak{d}') - 1) \deg(\mathfrak{p}).$$

- (2) *If \mathfrak{p} is fixed and \mathfrak{d}' varies, then*

$$\ln |\Phi_{J^\circ, \mathfrak{p}}| \sim 2h(\mathfrak{d}') \cdot c(\mathfrak{p}),$$

where $c(\mathfrak{p})$ is an explicit constant depending only on \mathfrak{p} , which can be estimated as $c(\mathfrak{p}) = \ln(|\mathfrak{p}| + \frac{1}{2}) + O(|\mathfrak{p}|^{-2} \deg(\mathfrak{p}))$.

Proof. It is known from the theory of automorphic forms that one of the roots of $P(x)$ is $\lambda_1 = |\mathfrak{p}| + 1$ and all other roots satisfy $|\lambda_i| \leq 2\sqrt{|\mathfrak{p}|}$, $2 \leq i \leq h(\mathfrak{d}')$. More precisely, the Brandt matrices form an algebra which is isomorphic to the algebra

of Hecke operators acting on the space spanned by one specific Eisenstein series and the “ \mathfrak{p} -new” part of Drinfeld automorphic cusp forms on the congruence subgroup $\Gamma_0(\mathfrak{d})$ of $\mathrm{GL}_2(A)$; cf. [21, §2]. Using this, one obtains the bound $|\lambda_i| \leq 2\sqrt{|\mathfrak{p}|}$ from the Ramanujan-Petersson conjecture over function fields. Part (1) then easily follows from Theorem 1.2.

If \mathfrak{p} is fixed and $\deg(\mathfrak{d}') \rightarrow \infty$, then the eigenvalues $\lambda_2, \dots, \lambda_{h(\mathfrak{d}')}$ become equidistributed in the interval $[-2\sqrt{|\mathfrak{p}|}, 2\sqrt{|\mathfrak{p}|}]$ with respect to a certain Sato-Tate measure. This follows from the equidistribution result of Nagoshi [12] for the eigenvalues of a fixed Hecke operator acting on the spaces of Drinfeld cusp forms as the level varies, and the observation of Serre [18, Thm. 2] that the same equidistribution persists if one restricts the Hecke operator to the new part of the space of cusp forms. Having this fact, one can argue as in [13, §4.3] to obtain part (2). \square

Acknowledgements. This work was carried out while I was visiting the Taida Institute for Mathematical Sciences in Taipei and the Max Planck Institute for Mathematics in Bonn. I thank these institutes for their hospitality and excellent working conditions. I am grateful to Fu-Tsun Wei for very useful discussions related to the topic of this paper.

2. GRAPHS AND LAPLACIANS

2.1. Graphs. A *graph* consists of a set of vertices $V(G)$, a set of oriented edges $E(G)$ and two maps




$$E(G) \rightarrow V(G) \times V(G), \quad e \mapsto (o(e), t(e))$$

and

$$E(G) \rightarrow E(G), \quad e \mapsto \bar{e}$$

such that $\bar{\bar{e}} = e$ and $t(\bar{e}) = o(e)$; cf. [19, p. 13]. For $e \in E(G)$, the edge \bar{e} is called the *inverse* of e , the vertex $o(e)$ (resp. $t(e)$) is called the *origin* (resp. *terminus*) of e . The vertices $o(e), t(e)$ are called the *extremities* of e . We say that two vertices are *adjacent* if they are the extremities of some edge. We say that G is *finite*, if it has finitely many vertices and edges. A *weighted graph* is a graph G equipped with two maps $w : E(G) \rightarrow \mathbb{Z}_{>0}$ and $w : V(G) \rightarrow \mathbb{Z}_{>0}$, called weight functions, such that $w(e) = w(\bar{e})$ for all $e \in E(G)$. We say that a weighed graph G is *N -regular* if there is a positive integer $N \geq 1$ such that for any vertex $v \in V(G)$ we have

$$\sum_{\substack{e \in E(G) \\ t(e)=v}} \frac{w(v)}{w(e)} = N.$$

Note that two vertices might be the extremities of multiple edges , i.e., there might be $e \neq e'$ with $o(e) = o(e')$ and $t(e) = t(e')$. Also, G might have loops , i.e., edges $e \in E(G)$ with $e \neq \bar{e}$ and $t(e) = o(e)$. Finally, as in [9], we do not exclude the case $e = \bar{e}$: .

We say that a group Γ acts on the weighted graph G if Γ acts on the sets $V(G)$ and $E(G)$ so that for any $\gamma \in \Gamma$, $v \in V(G)$, $e \in E(G)$ we have $\gamma(o(e)) = o(\gamma e)$, $\gamma \bar{e} = \gamma \bar{e}$, $w(\gamma e) = w(e)$, $w(\gamma v) = w(v)$. For a group Γ acting on G , we have a quotient graph

$\Gamma \setminus G$ and a natural mapping $j : G \rightarrow \Gamma \setminus G$ such that $V(\Gamma \setminus G) = \Gamma \setminus V(G)$ and $E(\Gamma \setminus G) = \Gamma \setminus E(G)$. Furthermore, if G is a weighted graph and the stabilizers

$$\Gamma_v := \{\gamma \in \Gamma \mid \gamma v = v\} \quad \text{and} \quad \Gamma_e := \{\gamma \in \Gamma \mid \gamma e = e\}$$

are finite for every $v \in V(G)$, $e \in E(G)$, then we make $\Gamma \setminus G$ into a weighted graph by putting $w(j(v)) = |\Gamma_v| \cdot w(v)$ and $w(j(e)) = |\Gamma_e| \cdot w(e)$.

Let K be a non-archimedean local field with finite residue field k . Let \mathcal{T} be the Bruhat-Tits tree associated with $\mathrm{SL}_2(K)$; cf. [19]. We make \mathcal{T} into a weighed graph by assigning the weight 1 to all its vertices and edges. Then \mathcal{T} is a $(|k| + 1)$ -regular tree. The group $\mathrm{PGL}_2(K)$ naturally acts on \mathcal{T} . Let Γ be a discrete subgroup of $\mathrm{PGL}_2(K)$ with compact quotient. Then the stabilizers in Γ of the vertices and edges of \mathcal{T} are finite, and, since Γ is cocompact in $\mathrm{PGL}_2(K)$, the quotient graph $\Gamma \setminus \mathcal{T}$ is finite; cf. [15, Lem. 5.1]. We make $\Gamma \setminus \mathcal{T}$ into a weighed graph as in the previous paragraph.

2.2. Laplacians. We assume from now on that G is finite and connected. Let $C_0(G, \mathbb{Q})$ be the vector space over \mathbb{Q} with basis $V(G)$. The (weighted) *Laplacian* is the linear transformation $\Delta : C_0(G, \mathbb{Q}) \rightarrow C_0(G, \mathbb{Q})$ defined by

$$\Delta(v) = \sum_{t(e)=v} \frac{w(v)}{w(e)}(v - o(e)).$$

It is well-known (and easy to prove) that the eigenvalues of Δ are non-negative real numbers, and 0 is an eigenvalue of Δ with multiplicity one; cf. [13, §2.1]. The weighted *adjacency operator* is the linear transformation $\delta : C_0(G, \mathbb{Q}) \rightarrow C_0(G, \mathbb{Q})$ defined by

$$\delta(v) = \sum_{t(e)=v} \frac{w(v)}{w(e)} o(e).$$

The pairing $E(G) \times E(G) \rightarrow \mathbb{Z}$:

$$(e, e') = \begin{cases} w(e) & \text{if } e' = e, \\ -w(e) & \text{if } e' = \bar{e}, \\ 0 & \text{otherwise,} \end{cases}$$

extends to a symmetric, bilinear, positive-definite pairing on the first simplicial homology group $H_1(G, \mathbb{Z})$, which is just a weighted version of the usual cycle pairing. Let $D(G)$ be the order of the cokernel of the map

$$\begin{aligned} H_1(G, \mathbb{Z}) &\longrightarrow \mathrm{Hom}(H_1(G, \mathbb{Z}), \mathbb{Z}) \\ \varphi &\mapsto (\varphi, *). \end{aligned}$$

For a graph G , let $E(G)^*$ be a subset of $E(G)$ with the following two properties:

- (i) if $e = \bar{e}$ then $e \notin E(G)^*$;
- (ii) if $e \neq \bar{e}$ then either e or \bar{e} , but not both, is in $E(G)^*$.

We define the *mass* of G as

$$m(G) := \sum_{v \in V(G)} w(v)^{-1}.$$

Theorem 2.1. *Assume G is finite with n vertices. Let*

$$0 = \lambda_1 < \lambda_2 \leq \dots \leq \lambda_n$$

be the eigenvalues of Δ . Then

$$D(G) = m(G)^{-1} \frac{\prod_{e \in E(G)^*} w(e)}{\prod_{v \in V(G)} w(v)} \prod_{i=2}^n \lambda_i.$$

Proof. See Theorem 3.1 in [13]. \square

The arithmetic significance of this theorem comes from the following. Let $\pi : X \rightarrow S$ be a semi-stable curve of genus $g \geq 1$ over a Dedekind scheme S of dimension 1. Recall that this means that π is a proper and flat morphism whose fibres $X_{\bar{s}}$ over the geometric points \bar{s} of S are reduced, connected curves of arithmetic genus g , and have only ordinary double points as singularities; cf. [11, Def. 10.3.14]. We assume that the generic fibre X_η of X is a smooth, projective, geometrically connected curve. Let $s \in S$ be a closed point, and $x \in X_s$ be a singular point. There exists a scheme S' , étale over S , such that any point $x' \in X' := X \times_S S'$ lying above x , belonging to a fiber $X'_{s'}$, is a split ordinary double point, and

$$\widehat{\mathcal{O}}_{X', x'} \cong \widehat{\mathcal{O}}_{S', s'} \llbracket u, v \rrbracket / (uv - c)$$

for some $c \in \mathcal{O}_{S', s'}$. Moreover, the valuation w_x of c for the normalized valuation of $\mathcal{O}_{S', s'}$ is independent of the choice of S', s' , and of x' . For the proof of these facts we refer to [11, Cor. 10.3.22].

One can associate a graph G_{X_s} to X_s , the so-called *dual graph* (cf. [11, p. 511]): Let k_s be the residue field at s . The vertices of G_{X_s} are the irreducible components of $X_s \times_{k_s} \overline{k_s}$, and each ordinary double point $x \in X_s$ defines an edge e_x whose extremities correspond to the irreducible components containing x (the two orientations of e_x correspond to a choice of one of the two branches passing through x as the origin of e_x). We assign the weight $w(e_x) = w_x$.

Theorem 2.2. *Let J be the Jacobian variety of X_η . Let \mathcal{J} be the Néron model of J over S . Let $\Phi_{J,s} := \mathcal{J}_{\bar{s}} / \mathcal{J}_{\bar{s}}^0$ be the group of connected components of \mathcal{J} at $s \in S$. Then $|\Phi_{J,s}| = D(G_{X_s})$.*

Proof. This follows from 11.5 and 12.10 in [6]. \square

Thus, Theorems 2.1 and 2.2 relate the order $|\Phi_{J,s}|$ to the eigenvalues of the weighted Laplacian of G_{X_s} .

Remark 2.3. Note that we have assigned weights only to the edges of G_{X_s} , and in fact, $|\Phi_{J,s}|$ depends only on the weights of the edges of G_{X_s} . To apply the formula in Theorem 2.1, one can assign arbitrary weights to the vertices of G_{X_s} , e.g., all 1. As we will see, in the situations where X arises from quaternion algebras, there will be a natural choice for the weights of vertices.

2.3. Involutions. Let G be a weighed N -regular finite connected graph. In addition, assume that G is bipartite, that is, $V(G)$ is a disjoint union $V(G) = O \sqcup I$ such that every edge $e \in E(G)$ has one of its extremities in O and the other in I ; in particular, G has no edges with $t(e) = o(e)$.

Let τ be a non-trivial involution acting on G (as a weighted graph) which interchanges O and I . Note that τ does not fix any vertices or edges, although it is possible that $\tau(e) = \bar{e}$. Let $G' := G / \langle \tau \rangle$ be the quotient of G under the action of τ .

This graph G' is a weighted graph in which $w(v') = w(v)$ and $w(e') = w(e)$, where $v \in V(G)$ is a preimage of $v' \in V(G')$ and $e \in E(G)$ is a preimage of $e' \in E(G')$. Note that G' might have edges e' with $\bar{e}' = e'$; this happens when $\tau(e) = \bar{e}$ for the preimage $e \in E(G)$ of e' . Also, G' might contain loops; this happens if there are two edges $e_1 \neq e_2$ in G such that $o(e_1) = o(e_2)$, $t(e_1) = t(e_2)$, and $\tau(e_1) = \bar{e}_2$. Finally, note that G' is again N -regular.

Let $\Delta : C_0(G, \mathbb{Q}) \rightarrow C_0(G, \mathbb{Q})$ be the weighted Laplacian of G :

$$\Delta(v) = Nv - \delta(v),$$

where δ is the weighted adjacency operator of G . Let Δ' be the weighted Laplacian of G' and δ' its weighted adjacency operator.

Lemma 2.4. *Let h be the number of vertices of G' . If $\lambda_1, \dots, \lambda_h$ are the eigenvalues of δ' , then $\pm\lambda_1, \dots, \pm\lambda_h$ are the eigenvalues of δ .*

Proof. It is clear that G has $2h$ vertices. It is also clear that δ and τ commute as linear operators on $C_0(G, \mathbb{R})$. Let v_1, \dots, v_h be the vertices in O , so that $\tau(v_1), \dots, \tau(v_h)$ are the vertices in I . Let $v'_1, \dots, v'_h \in V(G')$ be the images of v_1, \dots, v_h ; we take these vertices as a basis of $C_0(G', \mathbb{R})$. Let M be the matrix of δ' with respect to this basis. From what was said, it is easy to see that δ is represented by the $2h \times 2h$ matrix $\begin{pmatrix} 0 & M \\ M & 0 \end{pmatrix}$. Since

$$\begin{pmatrix} I_h & I_h \\ I_h & -I_h \end{pmatrix} \begin{pmatrix} 0 & M \\ M & 0 \end{pmatrix} \begin{pmatrix} I_h & I_h \\ I_h & -I_h \end{pmatrix}^{-1} = \begin{pmatrix} M & 0 \\ 0 & -M \end{pmatrix},$$

the claim follows. \square

The largest eigenvalue of δ' is N , which occurs with multiplicity 1.

Corollary 2.5. *Let $P(x)$ be the characteristic polynomial of δ' . Let $P'(x)$ be the derivative of $P(x)$. Enumerate the eigenvalues of δ' so that $N = \lambda_1 > \lambda_2 \geq \dots \geq \lambda_h$. The eigenvalues of Δ are $N \pm \lambda_i$, $1 \leq i \leq h$, and the product of non-zero eigenvalues of Δ is*

$$2N \prod_{i=2}^h (N + \lambda_i)(N - \lambda_i) = |P(-N)P'(N)|.$$

Proof. The first claim follows from Lemma 2.4. The last equality follows from

$$P'(N) = \prod_{i=2}^h (N - \lambda_i), \quad P(-N) = (-1)^h 2N \prod_{i=2}^h (N + \lambda_i).$$

\square

3. GRAPHS ARISING FROM QUATERNION ALGEBRAS

The set-up and notation in this section will be the same as in the introduction. In particular, \mathcal{B} is an indefinite quaternion algebra over F of discriminant \mathfrak{d} . We fix a prime $\mathfrak{p} \nmid \mathfrak{d}$, and denote by \mathcal{B}' the definite quaternion algebra of discriminant $\mathfrak{d}' := \mathfrak{d}/\mathfrak{p}$. Let $K := F_{\mathfrak{p}}$, and

$$\begin{aligned} \Gamma_0 &:= (\mathcal{M} \otimes A[\mathfrak{p}^{-1}])^{\times} / A[\mathfrak{p}^{-1}]^{\times}, \\ \Gamma_+ &:= \left\{ m \in (\mathcal{M} \otimes A[\mathfrak{p}^{-1}])^{\times} \mid \text{ord}_{\mathfrak{p}} \text{Nr}(m) \in 2\mathbb{Z} \right\} / A[\mathfrak{p}^{-1}]^{\times}. \end{aligned}$$

Note that Γ_+ is a normal subgroup of Γ_0 of index 2. By fixing an isomorphism $\mathcal{B}' \otimes_F K \cong \mathbb{M}_2(K)$, one can consider Γ_0 and Γ_+ as discrete cocompact subgroups of $\mathrm{PGL}_2(K)$. Let \mathcal{T} be the Bruhat-Tits tree associated with $\mathrm{SL}_2(K)$. As we have discussed in §2.1, the quotient graphs $G_0 := \Gamma_0 \backslash \mathcal{T}$ and $G_+ := \Gamma_+ \backslash \mathcal{T}$ are finite, weighted, $(|\mathfrak{p}| + 1)$ -regular graphs.

Let $V(\mathcal{T}) = V(\mathcal{T})_1 \sqcup V(\mathcal{T})_2$ be the disjoint union such that, for $v \in V(\mathcal{T})_i$ and $u \in V(\mathcal{T})_j$, the combinatorial distance between v and u is even if and only if $i = j$. Using the Corollary on page 75 in [19], it is easy to see that $\gamma V(\mathcal{T})_i = V(\mathcal{T})_i$ ($i = 1, 2$) for $\gamma \in \Gamma_+$, and $\gamma V(\mathcal{T})_1 = V(\mathcal{T})_2$ and $\gamma V(\mathcal{T})_2 = V(\mathcal{T})_1$ for $\gamma \in \Gamma_0 - \Gamma_+$. Thus, G_+ is bipartite. Fix some $\tau \in \Gamma_0 - \Gamma_+$. It induces an involution of G_+ which does not depend on the choice of τ . By abuse of notation, we will denote this involution by τ . The triple $\{G_+, \tau, G_0 = G_+/\langle \tau \rangle\}$ fits into the setting of §2.3.

As is explained in [9, p. 291], using Eichler's approximation theorem, one can identify $V(G_0) = \{v_1, \dots, v_{h(\mathfrak{d}')}\}$ with the set $\{\mathcal{M}_1, \dots, \mathcal{M}_{h(\mathfrak{d}')}\}$ so that $(\Gamma_0)_{v'_i} \cong \mathcal{M}_i^\times / A^\times$, where v'_i is a preimage of v_i in \mathcal{T} . In particular, the cardinality of $V(G_0)$ is the class number of \mathcal{B}' and $w(v_i) = w_i$, $1 \leq i \leq h(\mathfrak{d}')$. Moreover, by [9, p. 294] (see also [5, Prop. 4.4]), the weighted adjacency operator of G_0 is given by the transpose of the \mathfrak{p} -th Brandt matrix $B(\mathfrak{p})$ if we fix $v_1, \dots, v_{h(\mathfrak{d}')}$ as a basis of $C_0(G_0, \mathbb{R})$. (In [9], Kurihara considers only quaternion algebras over \mathbb{Q} , but his arguments apply also, essentially verbatim, to quaternion algebras over F .)

Lemma 3.1. *Let $v \in V(G_0)$ be a vertex of weight $q + 1$. For an edge $e \in E(G_0)$ with $t(e) = v$ we have either $w(e) = 1$ or $w(e) = q + 1$. Moreover,*

$$\#\{e \in E(G_0) \mid t(e) = v, w(e) = q + 1\} = 1 + (-1)^{\deg(\mathfrak{p})}.$$

Proof. Let $e' \in E(\mathcal{T})$ be a preimage of $e \in E(G_0)$. Similar to the case of vertices, one can show that the stabilizer $(\Gamma_0)_{e'}$ is isomorphic to $\mathcal{I}^\times / A^\times$ for a certain Eichler A -order \mathcal{I} in \mathcal{B}' . On the other hand, by [3, p. 383], \mathcal{I}^\times is isomorphic either to \mathbb{F}_q^\times or $\mathbb{F}_{q^2}^\times$. This implies the first claim.

The second claim can also be deduced from the results in [3], but a more direct argument is the following: Let $v' \in V(\mathcal{T})$ be a preimage of v . Through the identification discussed in the introduction, let \mathcal{M}' be the maximal order corresponding to v' . Then $\mathcal{M}'^\times \cong \mathbb{F}_{q^2}^\times$. Let γ be a generator of this group. We need to count the number of edges $e' \in E(\mathcal{T})$ such that $t(e') = v'$ and $\gamma e' = e'$. The set

$$\mathrm{Star}(v') = \{e' \in E(\mathcal{T}) \mid t(e') = v'\}$$

is in natural bijection with $\mathbb{P}^1(\mathbb{F}_{\mathfrak{p}})$, and through this bijection γ acts on $\mathrm{Star}(v')$ in the same way as a certain matrix in $\mathrm{GL}_2(\mathbb{F}_q)$ acts on $\mathbb{P}^1(\mathbb{F}_{\mathfrak{p}})$; cf. [9, p. 292]. The characteristic polynomial of γ , as a matrix in $\mathrm{GL}_2(\mathbb{F}_q)$, is an irreducible quadratic polynomial over \mathbb{F}_q . It is easy to check that such a matrix has two fixed points in $\mathbb{P}^1(\overline{\mathbb{F}}_q)$ and these points lie in $\mathbb{P}^1(\mathbb{F}_{q^2}) - \mathbb{P}^1(\mathbb{F}_q)$. Therefore, γ acting on $\mathbb{P}^1(\mathbb{F}_{\mathfrak{p}})$ has no fixed points if $\deg(\mathfrak{p})$ is odd, and has exactly two fixed points if $\deg(\mathfrak{p})$ is even. \square

Theorem 3.2. *The graph G_+ has the following properties:*

- (1) G_+ is a finite, weighted, $(|\mathfrak{p}| + 1)$ -regular graph.
- (2) $|V(G_+)| = 2h(\mathfrak{d}')$.
- (3) For $v \in V(G_+)$, we have $w(v) = 1$ or $q + 1$. The number of vertices of weight $q + 1$ is $2h_{q+1}$.

- (4) $\sum_{v \in V(G_+)} w(v)^{-1} = 2m(\mathfrak{d}')$.
- (5) For $e \in E(G_+)$, we have $w(e) = 1$ or $q + 1$. Modulo the orientation, the number of edges of G_+ of weight $q + 1$ is $h_{q+1} (1 + (-1)^{\deg(\mathfrak{p})})$.
- (6)

$$D(G_+) = \frac{|P(-|\mathfrak{p}| - 1) \cdot P'(|\mathfrak{p}| + 1)|}{2m(\mathfrak{d}')(q + 1)^{n(\mathfrak{d})}}.$$

Proof. We have $|V(G_+)| = 2|V(G_0)| = 2h(\mathfrak{d}')$, the weight of $v \in V(G_+)$ is the weight of its image in G_0 , and every $v \in V(G_0)$ has exactly two preimages in G_+ . This proves (1)-(4). Similarly, the weight of $e \in E(G_+)$ is equal to the weight of its image in G_0 . Therefore, by Lemma 3.1, for $e \in E(G_+)$ we have $w(e) = 1$ or $q + 1$. We can count the number of edges in G_0 of weight $q + 1$ using the same lemma. More precisely, for each of the h_{q+1} vertices of G_0 with weight $q + 1$ we get $(1 + (-1)^{\deg(\mathfrak{p})})$ edges e of weight $q + 1$ terminating at the given vertex. This way we count each edge with non-trivial weight twice if $e \neq \bar{e}$ (once from $t(e)$ and once from $t(\bar{e})$), and count the edge only once if $e = \bar{e}$. On the other hand, an edge of G_0 has two preimages in G_+ if $e \neq \bar{e}$, and one preimage if $e = \bar{e}$. Thus, modulo the orientation, the number of edges of G_+ of weight $q + 1$ is $h_{q+1} (1 + (-1)^{\deg(\mathfrak{p})})$. We proved that

$$\frac{\prod_{e \in E(G_+)} w(e)}{\prod_{v \in V(G_+)} w(v)} = \frac{(q + 1)^{h_{q+1}(1 + (-1)^{\deg(\mathfrak{p})})}}{(q + 1)^{2h_{q+1}}} = \frac{1}{(q + 1)^{n(\mathfrak{d})}}.$$

Now the formula in part (6) follows from Theorem 2.1 and Corollary 2.5. \square

As follows from the analogue of the Cherednik-Drinfeld uniformization for $X_K^\mathfrak{d}$, proven in this context by Hausberger [7], $X_K^\mathfrak{d}$ is a twisted Mumford curve. More precisely, if we denote by $\mathcal{O}_{\mathfrak{p}}^{(2)}$ the ring of integers of the quadratic unramified extension of K and denote by $\mathbb{F}_{\mathfrak{p}}^{(2)}$ the residue field of $\mathcal{O}_{\mathfrak{p}}^{(2)}$, then $X_K^\mathfrak{d}$ has a semi-stable model $X_{\mathcal{O}_{\mathfrak{p}}^{(2)}}^\mathfrak{d}$ over $\mathcal{O}_{\mathfrak{p}}^{(2)}$ such that the dual graph of $X_{\mathbb{F}_{\mathfrak{p}}^{(2)}}^\mathfrak{d}$ is the weighted graph G_+ . Therefore, using Theorem 2.2 and Theorem 3.2, we obtain Theorem 1.2.

4. SHIMURA CURVES OVER \mathbb{Q}

Let \mathcal{B} be an indefinite quaternion algebra over \mathbb{Q} of discriminant d . Let X^d be the corresponding Shimura curve over \mathbb{Q} parametrizing abelian surfaces equipped with an action of \mathcal{B} . The Jacobian variety J^d of X^d has bad reduction at the primes dividing d .

Fix a prime p dividing d . Let \mathcal{B}' be the rational definite quaternion algebra of discriminant $d' = d/p$. Fix a maximal \mathbb{Z} -order \mathcal{M} in \mathcal{B}' . Let $h(d')$ be the class number of \mathcal{B}' . Let $I_1, I_2, \dots, I_{h(d')}$ be representatives of the left ideal classes of \mathcal{M} , and let \mathcal{M}_i be the right order of I_i , $1 \leq i \leq h(d')$. Denote $w_i = \mathcal{M}_i^\times / \mathbb{Z}^\times$. We have Eichler's formula for the mass of \mathcal{B}' :

$$m(d') := \sum_{i=1}^{h(d')} w_i^{-1} = \frac{1}{12} \prod_{\substack{\ell | d' \\ \ell \text{ prime}}} (\ell - 1).$$

Let $B(p) \in \mathbb{M}_h(\mathbb{Z})$ be the p -th Brandt matrix for \mathcal{B}' , and $P(x)$ be the characteristic polynomial of $B(p)$. Write $(p+1) = \lambda_1 > \lambda_2 \geq \dots \geq \lambda_{h(d')}$ for the eigenvalues of $B(p)$.

Let

$$\Gamma_+ = \left\{ \gamma \in (\mathcal{M} \otimes \mathbb{Z}[p^{-1}])^\times \mid \text{ord}_p \text{Nr}(\gamma) \in 2\mathbb{Z} \right\} / \mathbb{Z}[p^{-1}]^\times.$$

The group Γ_+ acts on the Bruhat-Tits tree \mathcal{T} of $\text{SL}_2(\mathbb{Q}_p)$. The quotient $G_+ := \Gamma_+ \backslash \mathcal{T}_p$ is a finite weighted $(p+1)$ -regular bipartite graph.

The Cherednik-Drinfeld uniformization theorem for $X_{\mathbb{Q}_p}^d$ implies that this curve has a semi-stable model over \mathbb{Z}_p whose dual graph, as a weighted graph, is G_+ ; see [2] or [9]. Therefore, the order of the component group $\Phi_{J^d, p}$ of the Néron model of J^d at p is $|\Phi_{J^d, p}| = D(G_+)$. Now the same argument as in Section 3, with the same references to [9], produces the following formula:

$$|\Phi_{J^d, p}| = \frac{|P(-p-1)P'(p+1)|}{2m(d')} \frac{\prod_{e \in E(G_+)^*} w(e)}{\prod_{v \in V(G_+)} w(v)}.$$

The only small difference from the function field case is that now the weights $w(v)$, $w(e)$ for $v \in V(G_+)$ and $e \in E(G_+)$ are more varied, in the sense that $w(v)$ can be 1, 2, 3, 6, 12, and $w(e)$ can be 1, 2, 3. We consider three separate cases:

Case 1: $d' = 2$ (equiv. $d = 2p$). In this case, $h(d') = 1$, $m(d') = 1/12$, $P(x) = x - (p+1)$, and G_+ has two vertices of weight 12. By Proposition 4.2 in [9], the number of edges of weight 2 is $\frac{1}{2} \left(1 + \left(\frac{-4}{p} \right) \right)$, and the number of edges of weight 3 is $\left(1 + \left(\frac{-3}{p} \right) \right)$. (Here $\left(\frac{*}{p} \right)$ is the Legendre symbol.) Therefore

$$(4.1) \quad |\Phi_{J^d, p}| = (p+1)2^{\frac{1}{2}(-3+(\frac{-4}{p}))}3^{(\frac{-3}{p})}.$$

Case 2: $d' = 3$ (equiv. $d = 3p$). In this case, $h(d') = 1$, $m(d') = 1/6$, $P(x) = x - (p+1)$, and G_+ has two vertices of weight 6. By Proposition 4.2 in [9], the number of edges of weight 2 is $\left(1 + \left(\frac{-4}{p} \right) \right)$, and the number of edges of weight 3 is $\frac{1}{2} \left(1 + \left(\frac{-3}{p} \right) \right)$. Therefore

$$(4.2) \quad |\Phi_{J^d, p}| = (p+1)2^{(-1+(\frac{-4}{p}))}3^{\frac{1}{2}(-1+(\frac{-3}{p}))}.$$

Case 3: $d' \geq 5$. In this case, the vertices of G_+ have weights 1, 2, or 3. Moreover, by [9, p. 291], the number of vertices of weight 2 (resp. 3) is $2h_2$ (resp. $2h_3$), where

$$h_2 = \frac{1}{2} \prod_{\ell|d'} \left(1 - \left(\frac{-4}{\ell} \right) \right), \quad h_3 = \frac{1}{2} \prod_{\ell|d'} \left(1 - \left(\frac{-3}{\ell} \right) \right).$$

By [9, Prop. 4.2] and the argument in Section 3, the number of edges (modulo the orientation) of weight 2 (resp. 3) in G_+ is $h_2 \left(1 + \left(\frac{-4}{p} \right) \right)$ (resp. $h_3 \left(1 + \left(\frac{-3}{p} \right) \right)$). Therefore,

$$\frac{\prod_{e \in E(G_+)^*} w(e)}{\prod_{v \in V(G_+)} w(v)} = \frac{2^{h_2(1+(\frac{-4}{p}))}3^{h_3(1+(\frac{-3}{p}))}}{2^{2h_2}3^{2h_3}} = \frac{1}{2^{n_2}3^{n_3}},$$

where

$$n_2 = \frac{1}{2} \prod_{\ell|d} \left(1 - \left(\frac{-4}{\ell} \right) \right), \quad n_3 = \frac{1}{2} \prod_{\ell|d} \left(1 - \left(\frac{-3}{\ell} \right) \right).$$

We get

$$(4.3) \quad |\Phi_{J^d,p}| = \frac{|P(-p-1)P'(p+1)|}{2m(d')2^{n_2}3^{n_3}} = \frac{p+1}{m(d')2^{n_2}3^{n_3}} \prod_{i=2}^{h(d')} (p+1-\lambda_i)(p+1+\lambda_i).$$

Remark 4.1. Note that our formulas (4.1), (4.2) and (4.3) differ slightly from the formula in Theorem 2.3 of [8]. The formula in [8] is not correct as stated, which is a result of a subtraction mistake on line 2 of page 235 in [8].

REFERENCES

- [1] S. Bosch, W. Lütkebohmert, and M. Raynaud, *Néron models*, Springer-Verlag, 1990.
- [2] J.-F. Boutot and H. Carayol, *Uniformisation p -adique des courbes de Shimura: les théorèmes de Čerednik et de Drinfeld*, Astérisque (1991), no. 196-197, 45–158, Courbes modulaires et courbes de Shimura (Orsay, 1987/1988).
- [3] M. Denert and J. Van Geel, *The class number of hereditary orders in non-Eichler algebras over global function fields*, Math. Ann. **282** (1988), no. 3, 379–393.
- [4] E.-U. Gekeler and M. Reversat, *Jacobians of Drinfeld modular curves*, J. Reine Angew. Math. **476** (1996), 27–93.
- [5] B. Gross, *Heights and the special values of L -series*, Number theory (Montreal, Que., 1985), CMS Conf. Proc., vol. 7, Amer. Math. Soc., Providence, RI, 1987, pp. 115–187.
- [6] A. Grothendieck, *Modèles de Néron et monodromie*, SGA 7, Exposé IX, 1972.
- [7] T. Hausberger, *Uniformisation des variétés de Laumon-Rapoport-Stuhler et conjecture de Drinfeld-Carayol*, Ann. Inst. Fourier (Grenoble) **55** (2005), 1285–1371.
- [8] B. Jordan and R. Livné, *On the Néron model of Jacobians of Shimura curves*, Compositio Math. **60** (1986), no. 2, 227–236.
- [9] A. Kurihara, *On some examples of equations defining Shimura curves and the Mumford uniformization*, J. Fac. Sci. Univ. Tokyo **25** (1979), 277–300.
- [10] G. Laumon, M. Rapoport, and U. Stuhler, *\mathcal{D} -elliptic sheaves and the Langlands correspondence*, Invent. Math. **113** (1993), 217–338.
- [11] Q. Liu, *Algebraic geometry and arithmetic curves*, Oxford Graduate Texts in Mathematics, vol. 6, 2002.
- [12] H. Nagoshi, *The distribution of eigenvalues of arithmetic infinite graphs*, Forum Math. **14** (2002), no. 6, 807–829.
- [13] M. Papikian, *Graph Laplacians, component groups and Drinfeld modular curves*, Münster J. Math., to appear, available at <http://www.math.uni-muenster.de/mjm/>.
- [14] M. Papikian, *On Jacquet-Langlands isogeny over function fields*, J. Number Theory **131** (2011), no. 7, 1149–1175.
- [15] M. Papikian, *Local Diophantine properties of modular curves of \mathcal{D} -elliptic sheaves*, J. Reine Angew. Math. **664** (2012), 115–140.
- [16] M. Papikian and F.-T. Wei, *The Eisenstein ideal and Jacquet-Langlands isogeny over function fields*, Doc. Math. **20** (2015), 551–629.
- [17] A. Schweizer, *On the Drinfeld modular polynomial $\Phi_T(X, Y)$* , J. Number Theory **52** (1995), no. 1, 53–68.
- [18] J.-P. Serre, *Répartition asymptotique des valeurs propres de l'opérateur de Hecke T_p* , J. Amer. Math. Soc. **10** (1997), no. 1, 75–102.
- [19] J.-P. Serre, *Trees*, Springer Monographs in Mathematics, Springer-Verlag, 2003.
- [20] M.-F. Vignéras, *Arithmétique des algèbres de quaternions*, Lecture Notes in Mathematics, vol. 800, Springer, 1980.
- [21] F.-T. Wei and J. Yu, *Theta series and function field analogue of Gross formula*, Doc. Math. **16** (2011), 723–765.

DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16802, U.S.A.

E-mail address: papikian@psu.edu